

# Terrorism and Disaster

WHAT  
CLINICIANS  
NEED TO  
KNOW



## Staphylococcal Enterotoxin B

 RUSH UNIVERSITY  
MEDICAL CENTER



Sponsored for CME Credit by  
Rush University Medical Center

Release Date: April 1, 2005  
Expiration Date: March 31, 2007

# Terrorism and Disaster

WHAT  
CLINICIANS  
NEED TO  
KNOW

## SERIES EDITORS

**Rush University  
Medical Center  
Chicago, Illinois**

Stephanie R. Black, MD\*  
Assistant Professor of Medicine  
Section of Infectious Diseases  
Department of Internal Medicine

Daniel Levin, MD\*  
Assistant Professor  
General Psychiatry Residency Director  
Department of Psychiatry

Gillian S. Gibbs, MPH\*  
Project Coordinator  
Center of Excellence for Bioterrorism  
Preparedness

Linnea S. Hauge, PhD\*  
Educational Specialist  
Department of General Surgery

**AUTHORS  
Rush University  
Medical Center  
Chicago, Illinois**

Stephanie R. Black, MD\*  
Assistant Professor of Medicine  
Section of Infectious Diseases  
Department of Internal Medicine

Daniel Levin, MD\*  
Assistant Professor  
General Psychiatry Residency Director  
Department of Psychiatry

**Uniformed Services University  
Health Sciences  
Bethesda, Maryland**

David M. Benedek, MD, LTC, MC, USA  
Associate Professor of Psychiatry

Steven J. Durning, MD, Maj, USAF, MC\*  
Associate Professor of Medicine

Thomas A. Grieger, MD, CAPT, MC, USN\*  
Associate Professor of Psychiatry  
Associate Professor of Military &  
Emergency Medicine  
Assistant Chair of Psychiatry for Graduate  
& Continuing Education

Molly J. Hall, MD, Col, USAF, MC, FS\*  
Assistant Chair & Associate Professor  
Department of Psychiatry

Derrick Hamaoka, MD, Capt, USAF, MC, FS\*  
Director, Third Year Clerkship  
Instructor of Psychiatry

Paul A. Hemmer, MD, MPH, Lt Col, USAF, MC\*  
Associate Professor of Medicine

Benjamin W. Jordan, MD, CDR, MC, USNR, FS\*  
Assistant Professor of Psychiatry

James M. Madsen, MD, MPH, COL, MC-FS, USA\*  
Associate Professor of Preventive Medicine  
and Biometrics  
Scientific Advisor, Chemical Casualty Care  
Division, US Army Medical Research Institute  
of Clinical Defense (USAMRICD), APG-EA

Deborah Otori, MD, MPH, FACP, COL, MC, USA\*  
Associate Professor of Medicine

Michael J. Roy, MD, MPH, FACP, LTC, MC\*  
Associate Professor of Medicine  
Director, Division of Internal Medicine

Jamie Waselenko, MD, FACP\*\*  
Assistant Professor of Medicine  
Assistant Chief, Hematology/Oncology  
Walter Reed Army Medical Center  
Washington, DC

## Guest Faculty

Ronald E. Goans, PhD, MD, MPH\*  
Clinical Associate Professor  
Tulane University School of Public Health &  
Tropical Medicine  
New Orleans, LA

Sunita Hanjura, MD\*  
Rockville Internal Medicine Group  
Rockville, MD

Niranjan Kanesa-Thanan, MD, MTMH\*  
Director, Medical Affairs & Pharmacovigilance  
Acambis  
Cambridge, MA

Jennifer C. Thompson, MD, MPH, FACP\*  
Chief, Department of Clinical Investigation  
William Beaumont Army Medical Center  
El Paso, TX

## Faculty Disclosure Policy

It is the policy of the Rush University Medical Center Office of Continuing Medical Education to ensure that its CME activities are independent, free of commercial bias and beyond the control of persons or organizations with an economic interest in influencing the content of CME. Everyone who is in a position to control the content of an educational activity must disclose all relevant financial relationships with any commercial interest (including but not limited to pharmaceutical companies, biomedical device manufacturers, or other corporations whose products or services are related to the subject matter of the presentation topic) within the preceding 12 months. If there are relationships that create a conflict of interest, these must be resolved by the CME Course Director in consultation with the Office of Continuing Medical Education prior to the participation of the faculty member in the development or presentation of course content.

\* Faculty member has nothing to disclose.

\*\*Faculty disclosure: CBCE Speaker's Core for SuperGen.

# Staphylococcal Enterotoxin B

CASE AUTHOR: Paul A. Hemmer, MD, MPH, Lt Col, USAF, MC

## ACCREDITATION & DESIGNATION STATEMENT

Rush University Medical Center is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide continuing medical education for physicians.

Rush University Medical Center designates this educational activity for a maximum of 1 credit of category 1 credit toward the AMA Physician's Recognition Award. Each physician should claim only those credits that he/she actually spent in the activity.

This CME activity was planned and produced in accordance with the ACCME Essentials.

CME credits are available free of charge through March 2007.

## DISCLAIMER

This project was funded by the Metropolitan Chicago Healthcare Council (MCHC) through a grant from the Health Resources and Services Administration (HRSA).

The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as representing the opinion of Rush University Medical Center, the Department of the Army, Department of the Navy, Department of the Air Force, Department of Defense, MCHC or HRSA.

## FDA Approved Drug and Devices Assurance Statement

In accordance with requirements of the FDA, the audience is advised that information presented in this continuing medical education activity may contain references to unlabeled or unapproved uses of drugs or devices. Please refer to the FDA approved package insert for each drug/device for full prescribing/utilization information.

## INSTRUCTIONS

The questions that appear throughout this case are intended as a self-assessment tool. For each question, select or provide the answer that you think is most appropriate and compare your answers to the key at the back of this booklet. The correct answer and a discussion of the answer choices are included in the answer key.

Note: These self-assessment questions are not intended for CME credit. To apply for CME credit, you must complete the CME Test at the back of this booklet and submit it according to the directions provided.

In addition, a sign is provided in the back of this booklet for posting in your office or clinic. Complete the sign by adding your local health department's phone number.

*Design and layout © 2005 Rush University Medical Center. The text contained herein falls under the U.S. Copyright Act of 1976 as a "U.S. Government Work" and is therefore considered Public Domain Information, however Rush University Medical Center reserves the right to copyright the design and layout of that information.*

# Staphylococcal Enterotoxin B

CASE AUTHOR: Paul A. Hemmer, MD, MPH, Lt Col, USAF, MC

## INTENDED AUDIENCE

Internal medicine, family medicine, and emergency medicine physicians, and other clinicians who will provide evaluation and care in the aftermath of a terrorist attack or other public health disaster

## LEARNING OBJECTIVES

Upon completion of this case, participants will be able to:

- Discuss the initial presentation of acute gastrointestinal illness due to Staphylococcal enterotoxin B (SEB), and contrast this with other common agents that occur naturally or that may be the result of intentional poisoning.
- Describe the natural and intentional sources of transmission and spread of SEB and other common biologic agents.
- List treatment options for patients with SEB.
- Summarize information that patients need to know about toxin-mediated illness.

## CASE PRESENTATION

You receive a phone call that one of your patients, a 45-year-old male, has presented to the Emergency Department (ED) late in the morning with complaints of recurring episodes of vomiting over the past 3 hours. You are informed by the ED attending that your patient's symptoms began with a sudden onset of intense nausea, some abdominal cramping, followed quickly by repeated emesis. You are told that your patient believes he has vomited 10 times in the past 3 hours and cannot tolerate anything by mouth. He reports having had 1 loose bowel movement. He denies witnessing blood in his emesis or stool, and there has been no coffee ground emesis. There is no history of fever. He was well upon waking this morning and until the onset of symptoms. He has been attending a local political convention for the past 3 days. Shortly after his arrival in the ED, 3 fellow convention attendees present with similar symptoms.

---

## QUESTION 1

At this time, likely etiologies of your patient's symptoms could include:

- a. Ingestion of Staphylococcal enterotoxin B (SEB), or *Bacillus cereus*
  - b. Ingestion of *Bacillus anthracis*, or *Clostridium botulinum* toxin
  - c. Fish or shellfish poisoning
- 

**Reminder:** You can find the Answer Key & Discussion on page 9.

**COMMENT:** Acute gastrointestinal illness due to ingestion of contaminated food is a very common problem, with estimates as high as 81,000,000 cases annually.<sup>1</sup> **The diagnosis of foodborne disease should be considered when acute illness with gastrointestinal or neurologic symptoms affects 2 or more people who have shared a meal within the past 72 hours.**<sup>2</sup>

While such illness is commonly acquired naturally, intentional contamination of food and water supplies as a weapon of bioterrorism is an increasing area of concern. Many of the common bacterial agents that cause acute gastrointestinal disease are listed by the Centers for Disease Control and Prevention (CDC) as Category B biologic agents for bioterrorism.<sup>3</sup> These Category B agents include: Staphylococcal enterotoxin B (SEB), *Salmonella* species, *Shigella dysenteriae*, *Escherichia coli* O157:H7, *Vibrio cholerae*, and *Cryptosporidium parvum*. Furthermore, two Category A agents may also cause acute gastrointestinal illness if ingested: *Bacillus anthracis* (anthrax), and *Clostridium botulinum* toxin (botulism). Because use of such agents is likely to be covert, people affected will be unaware of their exposure and will first come to attention when they seek medical care.<sup>3</sup> Thus, familiarity with the basic presenting clinical features is essential for early recognition, diagnosis, and treatment of infection with these agents.

The initial presentation of your patient with an acute vomiting syndrome should raise the suspicion of acute foodborne disease, particularly in light of 3 fellow convention attendees presenting with similar symptoms. In the initial approach to such patients, key historical information would include the symptom complex and trying to identify the incubation period of the illness (see Table 1). Illness due to the ingestion of preformed bacterial toxin will have the shortest incubation period, often on the order of a few hours. The most common bacterial agents to cause such illness are *Staphylococcus aureus* and *Bacillus cereus*. Both are likely to present with symptoms of nausea, repeated episodes of vomiting, and abdominal cramps within 1-6 hours of ingesting contaminated food. Diarrhea may be present. *S. aureus* may elaborate five distinct heat-stable toxins (A, B, C, D, and E), and the clinical manifestations of illness are similar.<sup>2</sup> SEB has relevance as a bioterrorism agent as it has already been isolated and produced.<sup>6</sup> **Fever is an uncommon manifestation of gastrointestinal illness with *S. aureus*.**

## CDC CLASSIFICATION OF BIOTERRORISM AGENTS

### Category A Agents

- Can be easily disseminated or transmitted person-to-person
- Cause high mortality with a potential large public health impact
- Might cause panic and social disruption
- Require special attention for public health preparedness

### Category B Agents

- Moderately easy to disseminate
- Cause moderate morbidity and low mortality
- Require enhancements to CDC's diagnostic and surveillance abilities

**Table 1. Category A and B Bioterrorism Agents\* Capable of Causing Acute Gastrointestinal Illness**

Agent [CDC Category]	Incubation Period, Range	Clinical Manifestations
<i>Staphylococcus aureus</i> [B]	1-6 hours	Vomiting, nausea, crampy abdominal pain, diarrhea, fever (~25%)
<i>Bacillus cereus</i> [B]	1-16 hours	Vomiting, crampy abdominal pain, diarrhea (~33%) Emetic syndrome: nausea, vomiting, 1-6 hours Diarrheal syndrome: abdominal cramping and diarrhea, 8-16 hours
<i>Clostridium Botulinum</i> [A]	18-36 hours†	Vomiting and nausea (~50%), diarrhea (~25%), descending paralysis
<i>Bacillus anthracis</i> (anthrax) [A]	3-7 days‡	Gastrointestinal disease: Initial: vomiting, nausea, abdominal pain, fever Later: severe abdominal pain, bloody diarrhea, toxemia, and intestinal perforation Oropharyngeal disease: neck swelling, lymphadenitis
<i>Salmonella</i> species [B]	16-72 hours†	Diarrhea, crampy abdominal pain, nausea, vomiting, fever, headache, rarely bloody diarrhea
<i>Shigella dysenteriae</i> [B]	16-72 hours†	Crampy abdominal pain, fever, diarrhea, bloody diarrhea, headache, nausea, vomiting
<i>Escherichia coli</i> O157:H7 [B]	72-120 hours†	Crampy abdominal pain, bloody diarrhea, fever (rare unless complicated by hemolytic-uremic syndrome), thrombotic thrombocytopenic purpura (developing 5-10 days after diarrhea) †
<i>Vibrio cholerae</i> [B]	16-72 hours†	Profound watery diarrhea, dehydration, fever (rare)‡
<i>Cryptosporidium parvum</i> [B]	5-28 days§	Diarrhea, may also have low grade fever, malaise, nausea, vomiting

\* As defined by the Centers for Disease Control and Prevention<sup>3</sup>

† Data from Tauxe, Swerdlow, and Hughes.<sup>2</sup>

‡ Data from Seas and Gotuzzo.<sup>4</sup>

§ Mean incubation period is 7-10 days

|| Data from Unger.<sup>5</sup>

*Bacillus cereus* produces a toxin that may lead to a short-incubation syndrome or emetic syndrome with an incubation period of 1-6 hours, nausea, abdominal cramps, vomiting, and less commonly diarrhea. *B. cereus* may also produce a diarrheal syndrome characterized by a longer incubation period of 8-16 hours, diarrhea and abdominal cramps. Vomiting is uncommon with the diarrheal syndrome, and fever is uncommon with both the emetic and diarrheal syndromes.<sup>2</sup>

*Clostridium botulinum* (botulism) also produces illness through a preformed toxin. Initial manifestations may include nausea and vomiting in up to 50% of affected people.<sup>2</sup> However, the incubation period may be longer, up to 18-36 hours. It is also characterized by the onset of a descending paralysis, typically causing bulbar palsies before descending to involve the extremities and respiratory musculature.<sup>7</sup>

Ingestion of *Bacillus anthracis* (anthrax) leads to nausea, vomiting, abdominal pain, and fever approximately 1-7 days after ingestion.<sup>7</sup> Symptoms rapidly worsen to include bloody diarrhea, severe abdominal pain with peritoneal signs, and occasionally hematemesis. Ascites may develop, as may bacteremia, gastrointestinal hemorrhage, or intestinal perforation.<sup>7</sup> Ingestion may also cause an oropharyngeal disease with neck swelling, lymphadenitis, and pharyngeal ulcers with a pseudomembrane.<sup>7</sup>

The other Category B agents listed by the CDC<sup>3</sup> will also cause an acute gastrointestinal illness. However, as summarized in Table 1, they tend to have a longer incubation period and often have diarrhea and fever as prominent clinical manifestations. *Shigella* and *E. coli* O157:H7 are common causes of bloody diarrhea.

Briefly, ingestion of heavy metals (copper, zinc, tin, and cadmium) may cause nausea, vomiting, and abdominal cramps within one hour of ingestion.<sup>2</sup> The presence of paresthesias along with abdominal symptoms should prompt consideration of various types of fish or shellfish poisoning, or ingestion of monosodium glutamate or niacin.

Approximately 4 hours before presenting to the ED, your patient had breakfast at a special interest group meeting involving several hundred delegates. The food was served in a buffet. He recalled having eggs, bacon, coffee, milk, and a small cream-filled pastry. His symptoms began approximately 2 hours after eating. The 3 other patients reported they had also been present at the morning meetings and had eaten similar foods, including eggs, bacon, sausage, cereal, milk, coffee, pastries, and a series of cold cut meats.

On examination, the patient's vital signs included: Blood Pressure: 115/65, with no orthostatic changes; Pulse 90 and regular; Respirations 14; Temperature 37.2°C. He is in mild distress, holding his abdomen, and vomits once during the examination. His abdomen is soft, non-tender, and there are diffusely increased bowel sounds.

---

## QUESTION 2

Foods commonly associated with the “natural” transmission of SEB include all of the following except:

- a. Fried rice
- b. Cream-filled pastries
- c. Ham
- d. Poultry
- e. Potato salads

---

**COMMENT:** The additional history, as well as the corroboration with others who are ill, suggest a causative organism with a short incubation period — likely *S. aureus* or *B. cereus*. Staphylococcal food poisoning has been associated with foods with a high salt or sugar content<sup>7</sup> such as ham, pork, canned beef, egg and potato salads, and cream-filled pastries.<sup>2</sup> In some cases, a food handler with a purulent skin infection can be identified as the source, but this is often not the case.<sup>2</sup> *B. cereus* is classically associated with fried rice, where the rice has been cooked and then held warm for a prolonged time.<sup>2</sup> The most commonly associated foods with transmission of the other Category A and B agents are listed in Table 2.



**Table 2. Foods Commonly Associated with Natural Transmission of CDC Category A and B Bioterrorism Agents\***

Agent [CDC category]	Common food sources
<i>Bacillus anthracis</i> (anthrax) [A]	Contaminated beef
<i>Clostridium Botulinum</i> [A]	Improperly home-canned vegetables, fruit, fish; honey (infant botulism)
Salmonella species [B]	Beef, poultry, eggs, dairy, produce
<i>Shigella dysenteriae</i> [B]	Milk, salads (egg, potato, tuna, turkey), lettuce
<i>Staphylococcus aureus</i> [B]	Ham, Pork, Canned Beef, Egg and Potato salads, cream filled pastries
<i>Escherichia coli</i> O157:H7 [B]	Salads, beef
<i>Vibrio cholerae</i> [B]	Contaminated water, shellfish
<i>Cryptosporidium parvum</i> [B]	Contaminated water

\*Data from Tauxe, Swerdlow, and Hughes,<sup>2</sup> Centers for Disease Control and Prevention,<sup>3</sup> and Hamer and Gorbach.<sup>7</sup>

### QUESTION 3

SEB may be chosen as an agent of bioterrorism because:

- It can be an effective lethal agent.
- It can be an effective incapacitating agent.
- It is spread effectively from person to person.
- It has a prolonged incubation period.

**COMMENT:** The use of any of these bacterial agents as a bioterrorism weapon may nullify these “classic” associations with specific food groups. Specifically, if the agents are to be deliberately used, they could be disseminated onto the most readily available food products. Thus, distribution of the agent on food served to large numbers of people (meetings, troop meals, grocery stores) would have the greatest impact.<sup>8</sup> Although such deliberate use of these agents seems difficult to consider, recent history points to their successful use. In 1984, 751 people in The Dalles, Oregon became ill following ingestion of Salmonella species deliberately placed on food supplies at local restaurants by a local religious faction hoping to influence election results.<sup>9</sup> There was also a case reported in 1997, in which 12 laboratory employees became ill after ingesting food contaminated with Shigella species; the investigation failed to identify a responsible party.<sup>10</sup>

The patient’s laboratory report demonstrated the following: normal complete blood count. Chemistries: Na 143, K 3.5, Cl 100, CO2 27, BUN 15, Creatinine 1.2, Glucose 104 mg/dL. Stool negative for occult blood, no fecal leukocytes.

### QUESTION 4

In patients whose illness is due to ingestion of SEB, what is the most likely acid-base disturbance to develop?

- Metabolic acidosis
- Metabolic alkalosis
- Respiratory acidosis
- Respiratory alkalosis



**COMMENT:** Because vomiting tends to be the most common and recurrent symptom, patients may develop a metabolic alkalosis.<sup>7</sup> In severe cases, volume depletion and prostration may occur<sup>7</sup> and may result in hemoconcentration and elevation in the BUN.

The diagnostic approach to patients with acute gastrointestinal illness, particularly when it appears to be associated with an outbreak, is aimed at attempting to isolate the causative organism. In the case of SEB, the diagnosis is made primarily on clinical and epidemiological grounds, as it is difficult to test for the toxin.<sup>11</sup> The diagnosis might also be confirmed by isolating *S. aureus* from the vomitus or feces of ill people and matching it to a sufficient amount of organism isolated from contaminated food or food handler's lesion.<sup>2</sup>

In general, the specimens to collect for laboratory analysis would include feces, vomitus, serum, and blood.<sup>2</sup> Cultures and samples of the food, the food preparation areas, and food handlers might also be warranted. Finally, it is important to notify and work with the laboratory so that any special collection or laboratory techniques that may not be readily known by practitioners can be used to identify all possible organisms.<sup>2</sup>

The patient was hydrated with intravenous normal saline and reported feeling better. He was discharged home with instructions to return if he developed a fever, bloody diarrhea, or was unable to tolerate oral intake. Subsequent investigation recovered *S. aureus* from the breakfast buffets used at all of the interest group meetings. Contaminated foods included the eggs, breakfast meats, and pastries. In all, 850 people became ill, and all voting at the convention had to be postponed. Although no arrests were made, authorities received a message from a terrorist cell within the US claiming responsibility and warning that this successful act was just the first of many planned attempts to poison the US food supply.

---

## QUESTION 5

Reasonable treatment options for patients suspected of having illness due to ingestion of SEB would include all of the following except:

- a. Hydration
- b. Anti-emetics
- c. Anti-diarrheal
- d. Dicloxacillin

---

**COMMENT:** Patients frequently request antibiotics because they do not know that antibiotics are not an effective treatment for toxin-related illnesses, such as SEB. Patients should be educated about toxin-related illnesses and the appropriate treatment options. See page 11 of this case booklet for an example of a patient education handout on ingestion of toxins. *You may detach this page and freely photocopy it for your patients.*

It is important to note that SEB has been isolated and produced as a powder, making it possible to disseminate the agent via aerosol. Thus, it has the potential to infect many more people, even miles from the delivery point, depending on prevailing winds.<sup>8</sup> The incubation period following inhalation of SEB is 3-12 hours.<sup>8</sup> Initial symptoms include severe headache, chills, myalgias, non-productive cough, and a high fever up to 41.1° Celsius (106° Fahrenheit).<sup>8</sup> While fever is not a common symptom of gastrointestinal SEB, it is typical of inhalational SEB. Dyspnea and chest pain may be present. Although diarrhea is unusual following inhalation, it may develop if patients were to swallow, as well as inhale, the agent.<sup>8</sup> The fever may last up to one week, and the cough up to one month.<sup>8</sup>

Physical exam findings may be nonspecific, including high fever and crackles on lung exam, and the chest x-ray is often normal except in severe cases where there may be increased interstitial markings, pulmonary edema, or adult respiratory distress syndrome.<sup>6,8</sup> The differential diagnosis of inhaled SEB would include flu-like illnesses, but there would likely be far more people presenting for evaluation following SEB inhalation than for a naturally occurring epidemic.<sup>8</sup> SEB differs from anthrax or pneumonic plague in that it does not rapidly progress to death, and differs from tularemia and plague due to lack of infiltrates on chest x-ray.<sup>8</sup>

Treatment of inhaled SEB is supportive, including oxygenation, fluid replacement, and ventilator support, if necessary.<sup>6</sup> Antibiotics are not indicated. Recovery is the rule, although patients may be incapacitated for days or weeks.<sup>8</sup>

## CASE SUMMARY

A microorganism that could be effective as a bioterrorism weapon should be:

- Able to cause illness in most who are exposed to the agent.
- Easily disseminated.
- Capable of causing infection despite different environmental exposures (eg, being distributed on a variety of foods).
- Available for production in sufficient quantity.<sup>11</sup>

The goal of a bioterrorist is to cause fear, uncertainty, a sense of vulnerability, and lack of control. These goals may be met by using agents that kill. However, the goals may also be accomplished by incapacitating large numbers of people, and this may, at times, be more effective than a lethal agent. Microorganism agents are likely to be used in a covert manner, so that the initial presentation of the attack will be patients who are ill, and presenting to their physicians or other health care providers.<sup>3</sup> For example, many of the Category B agents listed by the CDC<sup>3</sup> are capable of causing acute gastrointestinal illness, and some have already been used in a limited way.<sup>9,10</sup>

SEB meets many of the criteria for use as a bioterrorism weapon. Its primary use would be as an incapacitating agent, rather than a lethal agent.<sup>6</sup> The toxin has been isolated and produced, and the toxin is heat-stable. It has a high attack rate — most who are exposed through ingestion will become clinically ill.<sup>7</sup> The presenting symptoms following ingestion are non-specific and include nausea, repeated emesis, abdominal cramps, followed by onset of diarrhea. Fever is an uncommon manifestation of gastrointestinal illness from SEB. The short incubation period from exposure to onset of symptoms is characteristic of ingesting a preformed toxin, but such a history may not be immediately evident. While specific foods have been associated with SEB and other agents,<sup>2,7</sup> if used as a bioterrorism agent, such associations may not be valid. Laboratory studies that may help distinguish SEB from other agents would include the presence of fecal leukocytes and bloody stools in patients with invasive organisms. Although the toxin is difficult to test for, bodily fluids (vomitus, stool, blood, serum) should be collected to test for a causative organism, and samples from suspected food should also be collected. Antibiotics have a limited role in treating acute gastrointestinal illness, and have no role in the treatment of SEB. Supportive care and hydration therapy are indicated for SEB.

While it may seem a remote possibility that biologic agents will be used intentionally to contaminate food or water supplies, health care providers must remain vigilant to this possibility. Early identification of a potential outbreak may help speed the identification of the causative organism, improve intervention, and limit the psychological

impact (eg, fear, sense of vulnerability) of such an event. **Therefore, it is imperative that health care providers contact their local health departments if a toxin-related illness, such as SEB, is suspected.**

## ANSWER KEY & DISCUSSION

### QUESTION 1

At this time, likely etiologies of your patient’s symptoms could include:

- a. Ingestion of Staphylococcal enterotoxin B (SEB), or *Bacillus cereus*
- b. Ingestion of *Bacillus anthracis*, or *Clostridium botulinum* toxin
- c. Fish or shellfish poisoning

**ANSWER:** The correct answer is a. Ingestion of preformed toxin such as Staphylococcal enterotoxin B (SEB) or *Bacillus cereus* leads to the abrupt onset of nausea, vomiting, and abdominal symptoms. *Bacillus anthracis* and *Clostridium botulinum* toxin also have abdominal symptoms, but they are associated with longer incubation periods and are associated with bloody diarrhea in the former and descending paralysis in the latter. While fish or shellfish poisoning is possible, the lack of paresthesias makes this less likely at this point.

### QUESTION 2

Foods commonly associated with the “natural” transmission of SEB include all of the following except:

- a. Fried rice
- b. Cream-filled pastries
- c. Ham
- d. Poultry
- e. Potato salads

**ANSWER:** The correct answer is a. *Bacillus cereus* is classically associated with ingestion of fried rice, where the rice has been cooked and then held warm for a prolonged time.<sup>2</sup> All of the other listed foods are associated with transmission of SEB; cream-filled pastries and potato salads are commonly thought of foods associated with SEB.

### QUESTION 3

SEB may be chosen as an agent of bioterrorism because:

- a. It can be an effective lethal agent.
- b. It can be an effective incapacitating agent.
- c. It is spread effectively from person to person.
- d. It has a prolonged incubation period.

**ANSWER:** The correct answer is b. SEB has a high attack rate — most who are exposed through ingestion will become clinically ill<sup>5</sup> and incapacitated. SEB is not generally a lethal agent nor is it effectively transmitted from person to person, and it has a short incubation period of 1-6 hours.

## QUESTION 4

In patients whose illness is due to ingestion of SEB, what is the most likely acid-base disturbance to develop?

- a. Metabolic acidosis
- b. Metabolic alkalosis
- c. Respiratory acidosis
- d. Respiratory alkalosis

**ANSWER:** The correct answer is b. There are no specific laboratory findings associated with gastrointestinal illness due to SEB. Because vomiting tends to be the most common and recurrent symptom, patients may develop a metabolic alkalosis.<sup>5</sup> In severe cases, volume depletion and prostration may occur<sup>5</sup> and one may see laboratory evidence such as hemoconcentration, and elevation in the BUN.

## QUESTION 5

Reasonable treatment options for patients suspected of having illness due to ingestion of SEB would include all of the following except:

- a. Hydration
- b. Anti-emetics
- c. Anti-diarrheal
- d. Dicloxacillin

**ANSWER:** The correct answer is d. Antibiotics are not indicated in the treatment of SEB or *B. cereus*. Antibiotics are lifesaving in the treatment of salmonellosis and typhoid fever, but should not be used in nontyphoidal salmonella infections.<sup>2</sup> Treatment of many cases of food or water borne illness is focused on supportive measures, including hydration and anti-emetics. While anti-diarrheal agents may be used for some patients, they are to be avoided in patients with high fever, bloody stools, or fecal leucocytes (signifying invasive infection).<sup>2</sup>

## PATIENT EDUCATION

Your acute illness of vomiting, nausea, and diarrhea is likely due to a chemical (a toxin) naturally produced by bacteria in the food that you ate.

For most people, the illness will last about 12 hours. You should drink liquids and eat what you can tolerate without vomiting. Although your illness is most likely caused by a bacterial toxin, antibiotics will not help the problem. Your body will be able to clear the toxin in a matter of hours.

You should contact your doctor or seek medical attention if any of the following occur:

1. High fever (greater than 101.5°F)
2. Dizziness when standing (this may mean you haven't been able to drink enough fluids).
3. You cannot eat or drink anything for more than 12-24 hours.
4. There is blood in your stool or bloody diarrhea.

As a general rule, good hand washing, especially after using the restroom, will help prevent the spread of gastrointestinal illness to other people.

*Patient education material developed by Paul A. Hemmer, MD, MPH, Lt Col, USAF, MC, Associate Professor of Medicine, Uniformed Services University of the Health Sciences, Bethesda, Maryland.*

## REFERENCES

1. Archer DL, Kvenberg JE. Incidence and cost of foodborne diarrheal disease in the United States. *J Food Protection*. 1985;48:887-894.
2. Tauxe RV, Swerdlow DL, Hughes JM. Foodborne disease. In: Mandell GL, Bennett JE, Dolin R, eds. *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases*. Vol.1. 5th ed. Philadelphia, Pa: Churchill Livingstone, Inc.; 2000:1150-1164.
3. Centers for Disease Control and Prevention. Biological and Chemical Terrorism: Strategic Plan for Preparedness and Response. Recommendations of the CDC Strategic Planning Workgroup. *MMWR*. 2000;49(No.RR-4):1-14.
4. Seas C, Gotuzzo E. *Vibrio Cholerae*. In: Mandell GL, Bennett JE, Dolin R, eds. *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases*. Vol 1. 5th ed. Philadelphia, Pa: Churchill Livingstone, Inc.; 2000:2266-2271.
5. Ungar B. *Cryptosporidium*. In: Mandell GL, Bennett JE, Dolin R, eds. *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases*. Vol 1. 5th ed. Philadelphia, Pa: Churchill Livingstone, Inc.; 2000:2903-2912.
6. Blazes DL, Lawler JV, Lazarus AA. When biotoxins are tools of terror. *Postgrad Med*. 2002;112:89-95.
7. Hamer DH, Gorbach SL. Infectious diarrhea and bacterial food poisoning. In: Feldman M, Friedman LS, Sleisenger MH, eds. *Sleisenger & Fordtran's Gastrointestinal and Liver Disease: Pathophysiology/Diagnosis/Management*. 7th ed. Philadelphia, Pa: Saunders; 2002:1864-1913.
8. Franz DR, Jahrling PB, Friedlander AM, et al. Clinical recognition and management of patients exposed to biological warfare agents. *JAMA*. 1997;278:399-411.
9. Torok TJ, Tauxe RV, Wise R, et al. A large community outbreak of salmonellosis caused by intentional contamination of restaurant salad bars. *JAMA*. 1999;278:389-95.
10. Kolavic SA, Kimura A, Simons SL, et al. An outbreak of *Shigella dysenteriae* type 2 among laboratory workers due to intentional food contamination. *JAMA*. 1997;278:396-398.
11. Moran GJ. Threats in bioterrorism II: CDC category B and C agents. *Emerg Med Clin North Am*. 2002;20:311-330.

## SUGGESTED READING

1. Blazes DL, Lawler JV, Lazarus AA. When biotoxins are tools of terror. *Postgrad Med*. 2002;112:89-95.
2. Moran GJ. Threats in bioterrorism II: CDC category B and C agents. *Emerg Med Clin North Am*. 2002;20:311-330.

# EVALUATION FORM

## TERRORISM AND DISASTER: WHAT CLINICIANS NEED TO KNOW

### Staphylococcal Enterotoxin B

#### Participant Information

Name/Degree _____		Practice setting: <input type="checkbox"/> Hospital/In-patient <input type="checkbox"/> Outpatient/Clinic <input type="checkbox"/> Other
Address _____		Email _____
City _____		Clinical Specialty _____
State _____	Zip _____	Signature _____
Telephone _____		Date _____

#### Instructions for Physicians Receiving Credit

The questions that follow may be used to obtain continuing medical education credit. To obtain 1 hour of Category 1 credit towards the AMA Physician’s Recognition Award, read this case study, which will take one hour of your time, circle the correct answer to each of the CME questions, complete the evaluation form, and return both the CME question page and the evaluation form via mail or fax to:

Rush University Medical Center  
 Office of Continuing Medical Education  
 600 South Paulina Street, Suite 433 AAF  
 Chicago, Illinois 60612  
 Fax: (312) 942-2000

#### Case Study Evaluation

Please rate this case study according to the following scale: 1=Very Poor 2=Poor 3=Fair 4=Good 5=Very Good 6=Excellent

1. Accredited CME activities must be “free of commercial bias for or against any product.” In this regard, how would you rate this activity? If you perceived any bias, please provide specific comments below. 1 2 3 4 5 6

\_\_\_\_\_

2. How well did the case study satisfy your purpose for reading it?

3. To what extent were the stated objectives of the case study achieved?

4. In general, was the case study well organized and presented?

5. To what extent has this CME activity improved your preparedness to recognize and care for victims of a terrorism attack or other public health disaster?

6. What was your overall rating of this case study?

7. I would recommend this case study to a colleague.  Yes  No

8. Based upon your review of this case, what specific action(s) could you take to enhance disaster preparedness in your workplace? Please estimate the probability that you will act on this item (0-100% where 100% = certainty)

A. \_\_\_\_\_ A. \_\_\_\_\_

B. \_\_\_\_\_ B. \_\_\_\_\_

C. \_\_\_\_\_ C. \_\_\_\_\_

Please check this box if you prefer not to be contacted for follow-up about the impact of this activity on your clinical practice.



## QUESTIONS FOR CONTINUING MEDICAL EDUCATION

- 1. The incubation period following ingestion of Staphylococcal enterotoxin B (SEB) is:**
  - a. 1-6 hours
  - b. 8-16 hours
  - c. 16-24 hours
  - d. > 24 hours
- 2. Which of the following are NOT common symptoms following ingestion of SEB?**
  - a. Nausea and dehydration
  - b. Vomiting and diarrhea
  - c. Fever and abdominal pain
- 3. Patients presumed to have a gastrointestinal illness due to *Staphylococcus aureus* enterotoxin should be advised to contact their doctor if they experience which of the following?**
  - a. Mild abdominal cramping
  - b. Blood in their stool
  - c. Vomiting that lasts more than 1 hour
- 4. Which of the following would be appropriate treatment options for a patient ill with gastrointestinal symptoms due to SEB?**
  - a. Antiemetics, hydration, antibiotics
  - b. Hydration, antipyretics, antibiotics, antiemetics
  - c. Antipyretics, antiemetics, hydration
- 5. How does inhalation of SEB differ from ingestion of SEB?**
  - a. Fever is more commonly associated with ingestion of SEB.
  - b. The primary treatment modality for inhalation SEB is antibiotics.
  - c. Inhalation SEB is considered to be lethal.
  - d. Inhalation SEB has the potential to be quickly disseminated to more people.

# NOTES

## NOTES

# Terrorism and Disaster

WHAT  
CLINICIANS  
NEED TO  
KNOW

Rush University Medical Center faculty, in collaboration with faculty from the Uniformed Services University of the Health Sciences (USUHS) authored a case series to provide continuing medical education (CME) for terrorism preparedness and other public health emergencies.

A series of 14 case studies was developed to provide innovative learning opportunities for health professionals to problem-solve issues related to terrorism or other public health emergencies. Due to the complicated and volatile nature of a terrorist event, the case studies were designed to expand outside the clinician-patient interaction and involve:

- deploying outside resources
- notifying appropriate officials
- coordinating a response team
- dealing with media and concerned public
- initiating emergency/disaster plans

Each case provides the CME user with decision-making challenges within his or her discipline, along with scenarios that address broader interdisciplinary issues. This interdisciplinary approach is particularly important in disaster preparedness, when health professionals will likely be called on to work outside their day-to-day experiences.

Authored by experts in the field, each self-paced case includes a thorough case history, questions to test your knowledge, a resource list of additional readings and relevant websites. One-hour CME and CEU credit is available for each case, following the successful completion of the CME questions included with each case. The CME and CEU credit is available free of charge through March 2007.

*The cases in the series include:*

## MEDICINE

*The medicine cases address recognition of the agent, diagnosis, treatment, and medical case management.*

- **Pneumonic Plague**
- **Radiation Attack**
- **Sarin**
- **Smallpox**
- **Staphylococcal Enterotoxin B**
- **Viral Hemorrhagic Fevers**

## PSYCHIATRY

*The psychiatry cases address issues of disaster psychiatry.*

- **Emergency Mental Health After a Suicide Bombing**
- **Psychiatric Sequelae in a Survivor of 9/11**
- **Psychosocial Management of a Radiation Attack**

## INTERDISCIPLINARY

*The interdisciplinary cases address basic medical management, general disaster planning, communicating with the media and concerned public, and psychosocial case management.*

- **Inhalational Anthrax**
- **Pulmonary Toxicants**
- **SARS**
- **Smallpox**
- **Viral Encephalitis**

For more information or to order your free copy of any of the cases in this series, please contact:

Office of Continuing  
Medical Education  
Rush University Medical Center  
Suite 433 AAF  
Chicago, Illinois 60612  
Telephone: (312) 942-7119  
Facsimile: (312) 942-2000  
E-mail: [cme\\_info@rush.edu](mailto:cme_info@rush.edu)

 RUSH UNIVERSITY  
MEDICAL CENTER

