Center for the Study of Traumatic Stress

CSTS | Department of Psychiatry | Uniformed Services University | 4301 Jones Bridge Road, Bethesda, MD 20814-4799 | www.CSTSonline.org

Managing the Stress of Children After a Crisis

If children have been directly exposed to death and destruction following a disaster, parents need to communicate with them through various stages. Immediately, physical safety and security takes priority.However, in the aftermath, it is difficult to predict if they will develop psychological problems.

Uniformed Services Univ<u>ersity</u>

The following management plan may help minimize later difficulties.

- Discuss your own reactions with another adult before talking with your children. Your response to a crisis will affect your child's response. Make time to be available for your child to talk or just spend time together.
- Listen to your child's stories. Every child has a different way of responding to trauma. It is not advisable to require the same response of everyone.
- Discuss the event in an open and honest manner with your children. Children might want to talk intermittently, and younger children might need concrete information to be repeated.
- Be available for your child.
- Engage your child in conversation of their choosing not necessarily about their feelings or the scene. Talking about the normal events of life is central to health.
- Maintain daily routines and familiar schedules to the

If children have been directly exposed to death and destruction following a disaster, parents need to communicate with them through various stages. extent possible. Attending school, established family time, and bedtimes stories can be reassuring and help children to feel safe and normal. Now is not the time to introduce new routines.

■ Limit the times of exposure

to television or other sources on information about the disaster and its victims.

- Increase your child's sense of control and mastery within the household — let them plan dinner or the evening's activities.
- Reassure your children that the disaster was not their fault in any way.
- Help older children and adolescents understand their behavior and set limits for them.
- Be tolerant and understanding if your children return to old behaviors. These may include a loss of toilet training or the inability to fall asleep alone.
- Help your child modulate the extent to which they personalize or identify with the victims or the situation. Remind your children that they are safe and with you.
- Provide concrete information to your child about the disaster event and how it is different than their current environment (if applicable). This will help decrease over-identifying with the victims.



Center for the Study of Traumatic Stress Department of Psychiatry Uniformed Services University 4301 Jones Bridge Road, Bethesda, MD 20814-4799 www.CSTSonline.org